


Quantum Field Theory Exercises week 10

Exercise 13: UV aspects of ϕ^3 -theory

Consider the scalar ϕ^3 -theory, which has the Lagrangian $\mathcal{L} = \frac{1}{2}(\partial_\mu \phi)(\partial^\mu \phi) - \frac{1}{2}m^2\phi^2 - \frac{\lambda}{3!}\phi^3$. The momentum-space Feynman rules for this theory are basically the same as the ones that we have derived for the scalar ϕ^4 -theory, except for the fact that the interaction vertex now involves three ϕ -particles instead of four!

- Which amputated diagram contributes to the one-loop corrections to the ϕ^3 interaction?
- Argue that this contribution is finite. Why was this to be expected based on the discussion on page 28 of the lecture notes?
- Give the analytical expressions for the one-loop diagrams



with the solid lines indicating the scalar bosons.

Note: you are not supposed to perform the full calculation here!

Remark: the second diagram is usually called a “tadpole diagram”. It is not 1-particle irreducible and, consequently, it is not part of the 1-particle irreducible self-energy of the scalar particle. The fact that this tadpole diagram is non-zero implies that the field $\hat{\phi}$ actually has a non-vanishing vev at one-loop order, i.e. $\langle \Omega | \hat{\phi}(x) | \Omega \rangle = v \neq 0$. For a correct particle interpretation the theory should be reformulated in terms of the field $\hat{\phi}'(x) = \hat{\phi}(x) - v$, which has a vanishing vev. This reformulation procedure that removes the tadpole diagram from the scalar self-energy is referred to in the literature as “tadpole renormalization”.

- Suppose we regularize the loop integrals by means of a UV cutoff Λ . Discuss how the one-loop diagrams in part (c) will depend on this cutoff.
- Can the Lagrangian parameter m be finite?
- Repeat the power-counting analysis on page 87 of the lecture notes for the ϕ^3 theory. Determine the dimensionalities of spacetime n for which the theory is renormalizable, superrenormalizable and nonrenormalizable.
- Compare this to the discussion on page 28 of the lecture notes.
Hint: for n spacetime dimensions the action is given by $S = \int d^n x \mathcal{L}$. Use this to determine the mass dimension of the coupling constant λ for n spacetime dimensions.
- Take spacetime to be four-dimensional and draw all superficially divergent 1-particle irreducible one-loop diagrams, indicating their expected dependence on the UV cutoff Λ .

Exercise 14: generators of the Lorentz group

In order to prepare for the discussion on higher-spin theories, please do parts (a) and (b) of this exercise.

- (a) Consider an infinitesimal rotation by an angle $\delta\alpha$ about the \vec{e}_n -axis. Under this rotation the spatial components of the position four-vector x^μ transform according to

$$\vec{x} \rightarrow \vec{x}' \approx \vec{x} + \delta\alpha \vec{e}_n \times \vec{x} \equiv \vec{x} + \delta\vec{\alpha} \times \vec{x}.$$

Rewrite this infinitesimal transformation in the form $x'^\rho = \Lambda^\rho_\sigma x^\sigma \approx (g^\rho_\sigma + \omega^\rho_\sigma)x^\sigma$. We now want to determine $\omega_{\rho\sigma}$, with both indices down. Show that the nonzero $\omega_{\rho\sigma}$ components are given by $\omega_{12} = -\omega_{21} = (\delta\alpha)^3$, $\omega_{23} = -\omega_{32} = (\delta\alpha)^1$ and $\omega_{31} = -\omega_{13} = (\delta\alpha)^2$.

- (b) Consider an infinitesimal boost with velocity $\delta\vec{v}$. Under this boost the position four-vector transforms according to:

$$x^0 \rightarrow x^{0'} \approx x^0 + \delta\vec{v} \cdot \vec{x} \quad \text{and} \quad \vec{x} \rightarrow \vec{x}' \approx \vec{x} + x^0 \delta\vec{v}.$$

Rewrite this infinitesimal transformation in the form $x'^\rho = \Lambda^\rho_\sigma x^\sigma \approx (g^\rho_\sigma + \omega^\rho_\sigma)x^\sigma$. Show that the nonzero $\omega_{\rho\sigma}$ components are given by $\omega_{0k} = -\omega_{k0} = (\delta v)^k$ for $k = 1, 2, 3$.

- (c) On page 11 of the lecture notes the following has been derived for infinitesimal Lorentz transformations:

$$\phi(x) \rightarrow \phi'(x) = \phi(\Lambda^{-1}x) \approx \left(1 - \frac{i}{2} \omega_{\mu\nu} J^{\mu\nu}\right) \phi(x),$$

where $J^{\mu\nu} = i(x^\mu \partial^\nu - x^\nu \partial^\mu)$ are the six generators of the infinitesimal Lorentz transformations. Show that these generators satisfy the fundamental commutation relations

$$[J^{\mu\nu}, J^{\rho\sigma}] = i(g^{\nu\rho} J^{\mu\sigma} - g^{\mu\rho} J^{\nu\sigma} - g^{\nu\sigma} J^{\mu\rho} + g^{\mu\sigma} J^{\nu\rho}).$$

- (d) Introduce four $n \times n$ matrices γ^μ that satisfy $\{\gamma^\mu, \gamma^\nu\} = \gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2g^{\mu\nu} I_n$, with I_n the $n \times n$ unit matrix. Prove that also $S^{\mu\nu} \equiv \frac{i}{4} [\gamma^\mu, \gamma^\nu]$ satisfy the fundamental commutation relations for generators of the infinitesimal Lorentz transformations.
- (e) The infinitesimal Lorentz transformations of four-vectors in parts (a) and (b) can be written as:

$$V^\alpha \rightarrow V'^\alpha \approx (g^\alpha_\beta + \omega^\alpha_\beta) V^\beta = \left(g^\alpha_\beta - \frac{i}{2} \omega_{\mu\nu} (J^{\mu\nu})^\alpha_\beta\right) V^\beta,$$

with $(J^{\mu\nu})^\alpha_\beta = i(g^{\mu\alpha} g^\nu_\beta - g^\mu_\beta g^{\nu\alpha})$. Guess what ... show that also these generators $J^{\mu\nu}$ satisfy the fundamental commutation relations for generators of the infinitesimal Lorentz transformations.