## Quantum Field Theory Exercises week 8

## Exercise 10 (continued)

Complete exercise 10 by deriving Newton's gravitational potential in part b.

## Exercise 11: decay rates, cross sections and reaction channels

Read §2.8.3 of the lecture notes about CM kinematics and consider the scalar Yukawa theory.

- (a) Calculate the lowest-order decay width  $\Gamma_{\phi \to \psi \bar{\psi}}$  in the rest frame of the decaying particle. For which masses is this decay possible?
- (b) Consider the process  $\psi \bar{\psi} \rightarrow \phi \phi$  in the centre-of-mass frame.
  - Which channels contribute to this process?
  - What do the corresponding diagrams say about the  $\theta$ -dependence?
  - For which centre-of-mass energies can this process occur?
  - Calculate the lowest-order differential cross section  $\left( d\sigma/d\Omega \right)_{CM}$ .
- (c) Consider the process  $\psi \bar{\psi} \rightarrow \psi \bar{\psi}$  in the centre-of-mass frame.
  - Which channels contribute to this process?
  - What are the ranges of the Mandelstam variables?
  - For which centre-of-mass energies can this process occur?
  - Calculate the lowest-order differential cross section.
  - Assume that m > 2M. Look at the energy dependence of both reaction channels separately. Do you notice something special in one of them?
  - Use the other reaction channel to determine whether the Yukawa interaction between  $\psi$ -particles and  $\bar{\psi}$ -particles is attractive or repulsive.

Hint: don't perform an explicit calculation, just use the analogy with the calculation that is presented on pages 52 and 53 of the lecture notes.