

### Solution 11:

- (a) Consider the decay in the rest frame of the decaying particle:  $k_A^\mu = (m, \vec{0})$ ,  $p_1^\mu = (E_1, \vec{p})$  and  $p_2^\mu = (E_2, -\vec{p})$ . First, note that  $E_1 = E_2 = \sqrt{M^2 + \vec{p}^2} = m/2$  and therefore  $|\vec{p}| = p = \frac{1}{2} \sqrt{m^2 - 4M^2} \geq 0$ , which implies the condition  $m \geq 2M$ .

According to exercise 9(b) the matrix element is  $i\mathcal{M}^{LO}(\phi \rightarrow \psi\bar{\psi}) = -ig$ . Employing page 61 of the lecture notes with  $E_{CM}$  replaced by  $m$  gives the decay width

$$\Gamma_{\phi \rightarrow \psi\bar{\psi}} = \frac{1}{2m} g^2 \int d\Pi_2 \stackrel{p.63}{=} \frac{g^2}{2m} \frac{p}{16\pi^2 m} \int d\Omega = \frac{g^2}{16\pi m} \sqrt{1 - 4M^2/m^2},$$

where  $\int d\Omega = 4\pi$  has been used.

- (b) Consider the lowest-order scattering process  $\psi(k_A)\bar{\psi}(k_B) \rightarrow \phi(p_1)\phi(p_2)$  in the CM frame:

$$k_A^\mu = (E_A, 0, 0, k), \quad k_B^\mu = (E_B, 0, 0, -k), \quad p_1^\mu = (E_1, \vec{p}) \quad \text{and} \quad p_2^\mu = (E_2, -\vec{p}),$$

with  $E_A = E_B = E_1 = E_2 = \frac{1}{2} E_{CM} = \frac{1}{2} \sqrt{s}$ .

This implies:

$$|\vec{k}| = k = \frac{1}{2} \sqrt{s - 4M^2} \geq 0 \quad \Rightarrow \quad E_{CM} = \sqrt{s} \geq 2M,$$

$$|\vec{p}| = p = \frac{1}{2} \sqrt{s - 4m^2} \geq 0 \quad \Rightarrow \quad E_{CM} = \sqrt{s} \geq 2m,$$

and  $\vec{k}_A \cdot \vec{p} = kp \cos \theta = \frac{1}{4} \sqrt{s - 4M^2} \sqrt{s - 4m^2} \cos \theta$ .

According to exercise 9(c) the matrix element  $i\mathcal{M}^{LO}(\psi(k_A)\bar{\psi}(k_B) \rightarrow \phi(p_1)\phi(p_2))$  reads

$$\text{t-channel diagram} + \text{u-channel diagram} = -ig^2 \left( \frac{1}{(k_A - p_1)^2 - M^2} + \frac{1}{(k_A - p_2)^2 - M^2} \right),$$

where the diagram on the left is the  $t$ -channel diagram and the one on the right the  $u$ -channel. Bearing in mind that  $t = (k_A - p_1)^2 = m^2 + M^2 - \frac{s}{2} \left( 1 - \sqrt{1 - 4M^2/s} \sqrt{1 - 4m^2/s} \cos \theta \right)$  and  $u = (k_A - p_2)^2 = m^2 + M^2 - \frac{s}{2} \left( 1 + \sqrt{1 - 4M^2/s} \sqrt{1 - 4m^2/s} \cos \theta \right)$ , the matrix element and therefore also the differential cross section are symmetric under  $\cos \theta \rightarrow -\cos \theta = \cos(\pi - \theta)$ . Furthermore, note that the usual infinitesimal imaginary parts have been skipped in the above-given propagators. Since  $t, u < M^2$ , these propagators can never become on-shell and as such the infinitesimal imaginary parts are obsolete.

The lowest order differential cross section then reads

$$\frac{d\sigma^{LO}}{d\Omega} \stackrel{p.64}{=} \frac{\sqrt{1 - 4m^2/s}}{64\pi^2 s \sqrt{1 - 4M^2/s}} \left| \mathcal{M}^{LO}(\psi(k_A)\bar{\psi}(k_B) \rightarrow \phi(p_1)\phi(p_2)) \right|^2 \Theta(\sqrt{s} - 2M) \Theta(\sqrt{s} - 2m).$$

(c) Consider the lowest-order scattering process  $\psi(k_A)\bar{\psi}(k_B) \rightarrow \psi(p_1)\bar{\psi}(p_2)$  in the CM frame:

$$k_A^\mu = (E_A, 0, 0, k), \quad k_B^\mu = (E_B, 0, 0, -k), \quad p_1^\mu = (E_1, \vec{p}) \quad \text{and} \quad p_2^\mu = (E_2, -\vec{p}),$$

with  $E_A = E_B = E_1 = E_2 = \frac{1}{2} E_{CM} = \frac{1}{2} \sqrt{s}$ .

This implies:

$$|\vec{k}| = k = |\vec{p}| = p = \frac{1}{2} \sqrt{s - 4M^2} \geq 0 \quad \Rightarrow \quad E_{CM} = \sqrt{s} \geq 2M,$$

and  $\vec{k}_A \cdot \vec{p} = kp \cos \theta = \frac{1}{4} (s - 4M^2) \cos \theta$ .

Therefore we obtain:  $s \geq 4M^2$ ,  $t = (k_A - p_1)^2 = -\frac{s}{2} (1 - 4M^2/s)(1 - \cos \theta) \leq 0$   
 and  $u = (k_A - p_2)^2 = -\frac{s}{2} (1 - 4M^2/s)(1 + \cos \theta) \leq 0$ .

According to exercise 9(c) the matrix element  $i\mathcal{M}^{LO}(\psi(k_A)\bar{\psi}(k_B) \rightarrow \psi(p_1)\bar{\psi}(p_2))$  reads

$$+ \quad = -ig^2 \left( \frac{1}{s - m^2 + i\epsilon} + \frac{1}{t - m^2} \right),$$

where the diagram on the left is the s-channel diagram and the one on the right the t-channel.

The lowest order differential cross section then reads


$$\frac{d\sigma^{LO}}{d\Omega} \stackrel{p.64}{=} \frac{\left| \mathcal{M}^{LO}(\psi(k_A)\bar{\psi}(k_B) \rightarrow \psi(p_1)\bar{\psi}(p_2)) \right|^2}{64\pi^2 s} \Theta(\sqrt{s} - 2M).$$


A remarkable feature of this differential cross section originates from the  $s$ -channel contribution: it has a singularity at  $s = m^2$ , i.e. when the intermediate  $\phi$ -particle goes on-shell. This can occur if  $m > 2M$ , corresponding to an unstable  $\phi$ -particle (see part a). For this reason we kept the  $i\epsilon$  imaginary part in the propagator. Actually, when treated correctly this instability adds a finite imaginary piece to the denominator, resulting in a (finite) resonance. The occurrence of such resonances allows to discover new particles! A beautiful example of a recent discovery via a resonance is the discovery of the Higgs boson in 2012 (see attached figure). In the '90s of the last century the resonance of the  $Z$ -boson was studied in great detail at LEP (CERN, Geneva) and SLC (SLAC, Stanford). For instance this has led to the conclusion that there are three generations of light neutrinos (see attached plots).

By the way, from  $i\mathcal{M}^{LO}(\psi(k_A)\bar{\psi}(k_B) \rightarrow \psi(p_1)\bar{\psi}(p_2))$  it trivially follows that (see also Ex. 10)

$$V_{\bar{\psi}\psi}(r) = V_{\psi\psi}(r) = -\frac{(g/2M)^2}{4\pi r} e^{-mr},$$

which is a consequence of the invariance of the scalar Yukawa theory under crossing.





Number of light neutrino generations:  $N_\nu = 2.984 \pm 0.008$

