

## Solution 7:

- (a) A Lagrangian typically consists of kinetic and mass terms (containing two fields) as well as interaction terms (containing three or more fields). Thus, in the scalar Yukawa theory the interaction term is given by

$$\mathcal{L}_{\text{int}} = -g\psi^*\phi\psi = -\mathcal{H}_{\text{int}}.$$

- (b) First we expand the exponential:

$$e^{-i \int d^4x \hat{\mathcal{H}}_{\text{int}}(x)} = \hat{1} - i \int d^4x \hat{\mathcal{H}}_{\text{int}}(x) + \mathcal{O}(g^2) = \hat{1} - ig \int d^4x \hat{\psi}_I^\dagger(x) \hat{\phi}_I(x) \hat{\psi}_I(x) + \mathcal{O}(g^2).$$

At order  $\mathcal{O}(g^0)$  one obtains

$$\langle 0 | T(\hat{\psi}_I^\dagger(x_1) \hat{\phi}_I(x_2) \hat{\psi}_I(x_3) \hat{1}) | 0 \rangle = 0,$$

since according to Wick's theorem only a fully contracted set of fields can give rise to a nonzero vacuum expectation value. Three fields can simply not be fully contracted as each Wick contraction involves a pair of fields.

At order  $\mathcal{O}(g)$  one has:

$$-ig \int d^4x \langle 0 | T(\hat{\psi}_I^\dagger(x_1) \hat{\phi}_I(x_2) \hat{\psi}_I(x_3) \hat{\psi}_I^\dagger(x) \hat{\phi}_I(x) \hat{\psi}_I(x)) | 0 \rangle.$$

Applying Wick's theorem one obtains only then a non-vanishing result if  $\hat{\phi}_I(x_2)$  is Wick-contracted with  $\hat{\phi}_I(x)$  and according to exercise 5(b)

1.  $\hat{\psi}_I^\dagger(x_1)$  is Wick-contracted with  $\hat{\psi}_I(x)$  and  $\hat{\psi}_I(x_3)$  with  $\hat{\psi}_I^\dagger(x)$ ,

$$-ig \int d^4x \langle 0 | \overbrace{\hat{\psi}_I^\dagger(x_1) \hat{\phi}_I(x_2) \hat{\psi}_I(x_3) \hat{\psi}_I^\dagger(x) \hat{\phi}_I(x) \hat{\psi}_I(x)}^{\text{Wick contractions}} | 0 \rangle$$

OR

2.  $\hat{\psi}_I^\dagger(x_1)$  is Wick-contracted with  $\hat{\psi}_I(x_3)$  and  $\hat{\psi}_I^\dagger(x)$  with  $\hat{\psi}_I(x)$ ,

$$-ig \int d^4x \langle 0 | \overbrace{\hat{\psi}_I^\dagger(x_1) \hat{\phi}_I(x_2) \hat{\psi}_I(x_3) \hat{\psi}_I^\dagger(x) \hat{\phi}_I(x) \hat{\psi}_I(x)}^{\text{Wick contractions}} | 0 \rangle.$$

Every Wick contraction gives a Feynman propagator, and therefore one obtains at order  $\mathcal{O}(g)$ :

$$\begin{aligned} & -ig \int d^4x \left[ D_F(x - x_1; M^2) D_F(x_2 - x; m^2) D_F(x_3 - x; M^2) \right. \\ & \quad \left. + D_F(x_3 - x_1; M^2) D_F(x_2 - x; m^2) D_F(x - x; M^2) \right], \end{aligned}$$

using the propagator conventions

$$\langle 0 | T(\hat{\psi}_I(x) \hat{\psi}_I^\dagger(y)) | 0 \rangle = \langle 0 | T(\hat{\psi}_I^\dagger(y) \hat{\psi}_I(x)) | 0 \rangle \equiv D_F(x - y; M^2)$$


and

$$\langle 0 | T(\hat{\phi}_I(x) \hat{\phi}_I(y)) | 0 \rangle \equiv D_F(x - y; m^2).$$

This has the structure

vertex  $(-ig \int d^4x) * \psi$ -propagator  $* \phi$ -propagator  $* \psi$ -propagator.

(c) The respective Feynman diagrams are:



The arrows in these Feynman diagrams represent the direction of particle-number flow. The operators  $\hat{\psi}_I(x_3)$  and  $\hat{\psi}_I(x)$  correspond to antiparticle creation or particle annihilation at the spacetime points  $x_3$  and  $x$ : hence, the arrows flow into these external/internal points of the Feynman diagrams. The operators  $\hat{\psi}_I^\dagger(x_1)$  and  $\hat{\psi}_I^\dagger(x)$  correspond to particle creation or antiparticle annihilation at the spacetime points  $x_1$  and  $x$ : hence, the arrows flow out of these external/internal points of the Feynman diagrams.

(d) Coordinate-space Feynman rules for Green's functions in the scalar Yukawa theory :

1. For each  $\phi$ -propagator  $\begin{array}{c} x_1 \quad x_2 \\ \bullet \text{-----} \bullet \end{array}$  insert  $D_F(x_1 - x_2; m^2)$

and for each  $\psi$ -propagator  $\begin{array}{c} x_1 \quad x_2 \\ \bullet \longleftarrow \bullet \end{array}$  insert  $D_F(x_1 - x_2; M^2)$ .

2. For each vertex  $\begin{array}{c} \diagup \bullet \text{----} \\ \diagdown \end{array} \begin{array}{c} x \\ \bullet \end{array}$  insert  $(-ig) \int d^4x$ .






3. For each external point  $\begin{array}{c} x \\ \bullet \text{---} \end{array}$  or  $\begin{array}{c} x \\ \bullet \text{--} \end{array}$  insert 1.

No symmetry factors are needed because all fields involved in the interaction are different.

(e) Translation of these Feynman rules to momentum space :

1. For each  $\phi$ -propagator  $\begin{array}{c} q \\ \bullet \text{---} \bullet \end{array}$  insert  $\frac{i}{q^2 - m^2 + i\epsilon}$

and for each  $\psi$ -propagator  $\begin{array}{c} q \\ \bullet \longrightarrow \bullet \end{array}$  insert  $\frac{i}{q^2 - M^2 + i\epsilon}$ .

2. For each vertex  insert  $-ig$ .
3. For each external point  or  insert  $e^{-iq \cdot x}$   
 and for each external point  or  insert  $e^{+iq \cdot x}$ .
4. Impose energy-momentum conservation at each vertex by fixing one of the momenta.
5. Integrate over each undetermined loop momentum  $p_j$ :  $\int \frac{d^4 p_j}{(2\pi)^4}$ .

### Solution 8:

Every interaction involves a pair  $\hat{\psi}^\dagger(z)\hat{\psi}(z)$ , and consequently for every in-flowing line there is an outflowing line. Since  $\psi$ -particles and  $\bar{\psi}$ -antiparticles have opposite particle number (which is a charge-like quantum number),  $\hat{\psi}_I(x)$  annihilates the charge associated with a  $\psi$ -particle at spacetime point  $x$ , whereas  $\hat{\psi}_I^\dagger(x)$  creates the same charge. As the charge of a  $\phi$ -particle is zero, because it has no  $\psi$ -particle number, the “total charge” = “number of  $\psi$ -particles – number of  $\bar{\psi}$ -antiparticles” is conserved at each interaction vertex. Furthermore, a propagator

$$\langle 0 | T(\hat{\psi}_I(x)\hat{\psi}_I^\dagger(y)) | 0 \rangle = \langle 0 | T(\hat{\psi}_I^\dagger(y)\hat{\psi}_I(x)) | 0 \rangle$$

creates the charge associated with a  $\psi$ -particle at spacetime point  $y$  and annihilates it at spacetime point  $x$ . So, the  $\psi$ -charge flows continuously from  $y$  to  $x$  in such a propagator, as indicated by the arrow convention. This would not be the case for the propagators  $\langle 0 | T(\hat{\psi}_I(x)\hat{\psi}_I(y)) | 0 \rangle$  and  $\langle 0 | T(\hat{\psi}_I^\dagger(x)\hat{\psi}_I^\dagger(y)) | 0 \rangle$ , which would lead to a clash of arrows and consequently to a violation of particle-number conservation. However, we have seen in exercise 5(b) that these propagators are zero and hence forbidden.

*In conclusion: particle-number conservation causes the arrows in a Feynman diagram to link up and form a continuous flow.*